


# CHEMISTRY



## WORKSHEET-10



**STP**

A PROJECT BY PUNJAB GROUP

**Worksheet – 10**  
**(A. Physical Chemistry)**  
**Fundamental Concepts**

**Q.1** Avogadro's number represents the number of:

- A) Atoms in 1g of helium gas
- B) Atoms in 24g of Mg
- C) Molecules in 35.5g of chlorine gas
- D) Electrons needed to deposit 24g Mg

**Q.2** Which one of the following terms is not used for ionic compounds?

- A) Formula unit
- B) Empirical formula
- C) Molecular formula
- D) Formula mass

**Q.3** 98g  $\text{H}_2\text{SO}_4$  contains number of moles of ions:

- A) 4.0 moles of ions
- B) 1 mole of ions
- C) 2 moles of ions
- D) 3.0 moles of ions

**Q.4** Cationic molecular ions are produced by:

- A) Radio waves
- B)  $\alpha$ -rays
- C) Beam of electrons
- D) Both B and C

**Q.5** Isotopes differ in:

- A) Properties which depend upon mass
- B) Arrangement of electrons in orbitals
- C) Chemical properties
- D) The extent to which they may be affected by electromagnetic field

**Q.6** A limiting reactant is the one which:

- A) Is taken in lesser quantity in grams as compared to other reactants
- B) Is taken in lesser quantity in volume as compared to the other reactants
- C) Gives the maximum amount of the product which is required
- D) Gives the minimum amount of the product under consideration

USE THIS SPACE FOR  
SCRATCH WORK

**Q.7** Symbol indicates not only the name of elements but also represents all of the following EXCEPT:

- A) One atom of an element
- B) Number of parts by mass of an element
- C) 1 gram atom of an element
- D) 1 amu

**Q.8** Which of the following is not mono-isotopic element?

- A) F
- B) Cl
- C) Au
- D) As

**Q.9** Which of the following statements is incorrect?

- A) Formation of uni-negative ion is exothermic
- B) Number of positive ions having group of atoms is less than number of negative ions having group of atoms
- C) X – rays and beam of electrons are used to produce positive ions of Ne
- D) Number of cationic molecular ions is less than number of anionic molecular ions and they are more stable

**Q.10** What volume of oxygen gas is required for the complete combustion of  $5\text{cm}^3$  of ethyne ( $\text{C}_2\text{H}_2$ )?

- A)  $12.5\text{cm}^3$
- B)  $13.0\text{cm}^3$
- C)  $13.5\text{cm}^3$
- D)  $14.0\text{cm}^3$

**Q.11** The relative atomic mass of boron, which consists of isotopes  $^{10}_5\text{B}$  and  $^{11}_5\text{B}$  is 10.8amu. What is the percentage of  $^{10}_5\text{B}$  atoms in the isotopic mixture?

- A) 0.8%
- B) 20%
- C) 8.0%
- D) 80%

**Q.12** How many carbon atoms are present in 34.2g of sucrose ( $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ )  $M_r = 342$ ?

- A)  $6.0 \times 10^{22}$
- B)  $3.6 \times 10^{25}$
- C)  $7.2 \times 10^{23}$
- D)  $3.6 \times 10^{24}$

**Q.13** What is the number of molecules in  $1000\text{cm}^3$  of nitrogen gas under room conditions?

- A)  $2.5 \times 10^{22}$
- B)  $3.5 \times 10^{22}$
- C)  $4.0 \times 10^{23}$
- D)  $4.5 \times 10^{26}$

USE THIS SPACE FOR  
SCRATCH WORK

**Q.14 Efficiency of a chemical reaction depends upon:**

- A) Greater amount of product
- B) Greater amount of reactant
- C) Less amount of product
- D) Less amount of reactant

**Q.15 How many total number of atoms are present in 49.0g of sulphuric acid (H<sub>2</sub>SO<sub>4</sub>)?**

- A)  $7 \times 3 \times 10^{23}$
- B)  $7 \times 8 \times 10^{23}$
- C)  $5 \times 6 \times 10^{23}$
- D)  $6 \times 6 \times 10^{23}$

**Q.16 An organic compound has empirical formula CH<sub>2</sub>O. If molar mass of the compound is 90 grams, then molecular formula of this organic compound would be (Ar of C = 12, H = 1.008 and O = 16):**

- A) C<sub>6</sub>H<sub>6</sub>O<sub>2</sub>
- B) C<sub>3</sub>H<sub>3</sub>O
- C) C<sub>9</sub>H<sub>9</sub>O<sub>3</sub>
- D) C<sub>3</sub>H<sub>6</sub>O<sub>3</sub>

**Q.17 How many bromine (Br) atoms are in 3 moles of bromine (Br) element?**

- A)  $3 \times 6.022 \times 10^{23}$  atoms
- B)  $79 \times 3 \times 6 \times 10^{23}$  atoms
- C)  $81 \times 3 \times 10^{23}$  atoms
- D)  $3 \times 6.022 \times 10^{23}$  atoms

**Q.18 Carbon dioxide (CO<sub>2</sub>) gas produced during combustion analysis of given organic compound is absorbed in 50% of KOH solution. It is a:**

- A) Chemical change only
- B) Physical change only
- C) May be physical or chemical change
- D) Neither physical nor chemical change

**Q.19 In the experimental determination of the percentage of carbon and hydrogen in an organic compound, water is absorbed by:**

- A) KOH
- B) MgCl<sub>2</sub>
- C) K<sub>2</sub>SO<sub>4</sub>
- D) Mg(ClO<sub>4</sub>)<sub>2</sub>

**Q.20 12g of magnesium (Mg) reacts with dilute sulphuric acid (H<sub>2</sub>SO<sub>4</sub>) to produce hydrogen (H<sub>2</sub>) gas. The amount of hydrogen (H<sub>2</sub>) gas produced is:**

- A) 4g
- B) 3g
- C) 2g
- D) 1g

USE THIS SPACE FOR  
SCRATCH WORK



**Q.25** All of the following terms are matched correctly w.r.t their definition EXCEPT:

**USE THIS SPACE FOR  
SCRATCH WORK**

Options	Term	Definition
A)	Relative atomic mass	It is the mass of one atom of an element as compared to the mass of an atom of carbon taken as 12
B)	Relative formula mass	It is sum of relative atomic mass of atoms of one formula unit of an ionic compound
C)	Relative molecular mass	It is the sum of relative atomic mass of atoms of one molecule of a covalent compound
D)	Mass number	It is sum of proton and neutrino

**Q.26** Identify the incorrect statement about yield:

A) Actual yield is less than theoretical yield

B) Percentage yield =  $\frac{\text{actual yield}}{\text{theoretical yield}} \times 100$

C) Experimental error does not affect actual yield

D) Efficiency of a chemical reaction depends on the amount of product

**Q.27** Isotopes of an element have all of the following different properties EXCEPT

A) They have different chemical properties

B) They have difference mass number

C) They have different number of neutrons

D) They have different half life

**Q.28** The combustion analysis of an organic compound shows 60% carbon, 8% hydrogen and 32% oxygen. If the molecular mass of the given organic compound is 200, then the molecular formula of the organic compound is (Ar of C = 12 amu, H = 1 amu and O = 16 amu):

A)  $C_{10}H_{16}O_4$

C)  $C_{10}H_{14}O_4$


B)  $C_8H_{16}O_4$

D)  $C_5H_8O_2$

**Q.29** Ascorbic acid (vitamin C) contains 48% carbon, 4% hydrogen and 48% oxygen. Which of the following is empirical formula of ascorbic acid?

- A)  $C_2H_4O_3$                       C)  $C_2H_2O_3$   
B)  $CH_2O$                          D)  $C_4H_4O_3$

**Q.30** Calcium reacts with excess oxygen to form calcium oxide (CaO) as shown in the equation:



The maximum mass of CaO formed when 4.0g of calcium is burnt in excess oxygen is ( $A_r$  values Ca = 40amu, O = 16amu):

- A) 3.6g                                C) 2.6g  
B) 5.6g                                D) 4.6g

**Q.31** If we know the mass of one substance, we can calculate the volume of other substance and vice versa with the help of a balanced chemical equation, which is called:

- A) Mass-mass relationship  
B) Mass-volume relationship  
C) Mole-volume relationship  
D) Mass-mole relationship

**Q.32** By using the value of Avogadro's number ( $N_A = 6.0 \times 10^{23}$ )mol<sup>-1</sup>, calculate the total number of atoms in 7.1g of Cl-element ( $A_r$  value Cl = 35.5):

- A)  $1.2 \times 10^{23}$  Cl-atoms            C)  $1.0 \times 10^{23}$  Cl-atoms  
B)  $1.6 \times 10^{23}$  Cl-atoms            D)  $1.5 \times 10^{23}$  Cl-atoms

**Q.33** Which one of the followings has same number of molecules as present in 11g of CO<sub>2</sub>?

- A) 4g of O<sub>2</sub>                            C) 4g of O  
B) 4.5g of H<sub>2</sub>O                        D)  $\frac{1}{4}$  moles of NaCl

**Q.34** 28g of N<sub>2</sub> gas at STP will occupy the volume of:

- A) 22.41dm<sup>3</sup>                            C) 44.82cm<sup>3</sup>  
B) 44.82dm<sup>3</sup>                            D) 2.241dm<sup>3</sup>

**USE THIS SPACE FOR  
SCRATCH WORK**

**Q.35** Which statements about a 12.0 g sample of  $^{12}\text{C}$  are correct?

- A) The number of atoms is  $6.02 \times 10^{23}$   
 B) The number of atoms is the same as the number of atoms in 4.0 g of  $^4\text{He}$ .  
 C) The number of atoms is the same as the number of atoms in 2.0 g of  $^1\text{H}_2$   
 D) Both A and B

**Q.36** A gaseous organic compounds, X, was burnt in an excess of oxygen. A  $0.112 \text{ dm}^3$  sample of X, measured at S.T.P., produced 0.88 g of carbon dioxide.

How many carbon atoms are there in one molecule of X?

- A) 1  
 B) 2  
 C) 3  
 D) 4

**Q.37** For complete oxidation, 1 mol of an organic compound requires 3 mol of oxygen gas. What could be the formula of the compound?


- A)  $\text{CH}_3\text{CHO}$   
 B)  $\text{CH}_3\text{CH}_2\text{OH}$   
 C)  $\text{CH}_3\text{CH}_3$   
 D)  $\text{CH}_3\text{CO}_2\text{H}$

**Q.38** A substance X was found to contain 72% carbon, 12% hydrogen and 16% oxygen.

What is the empirical formula of X?

- A)  $\text{C}_2\text{H}_4\text{O}$   
 B)  $\text{C}_3\text{H}_6\text{O}$   
 C)  $\text{C}_6\text{H}_{12}\text{O}$   
 D)  $\text{C}_6\text{H}_{12}\text{O}_2$

**Q.39** The relative atomic mass of chlorine is 35.5. which one of the following is the mass spectrogram of chlorine?



**USE THIS SPACE FOR  
SCRATCH WORK**



- Q.40  $10^{-3}$  moles of  $\text{MgSO}_4$  has the mass  
(Ar, Mg = 24 = 32 O = 16)
- A) 0.11g                      C) 0.13g  
B) 0.12g                      D) 0.14g

USE THIS SPACE FOR  
SCRATCH WORK

STEP ENTRY TEST 2021

**ANSWER KEY (Worksheet – 10)**

1	B	11	B	21	C	31	B
2	C	12	C	22	B	32	A
3	D	13	A	23	C	33	B
4	D	14	A	24	B	34	A
5	A	15	A	25	D	35	D
6	D	16	D	26	C	36	D
7	D	17	D	27	A	37	B
8	B	18	A	28	A	38	C
9	D	19	D	29	D	39	D
10	A	20	D	30	B	40	B

**ANSWERS EXPLAINED**

**Q.1 (B)** The number of particles present in one mole of a substance is called **Avogadro's number**.

( $N_A = 6.022 \times 10^{23}$ ) Statement (B) fulfills the condition of  $N_A$  such as. 24g of Mg = 1 mole

Molar mass of Mg = 24g  
=  $6.022 \times 10^{23}$  Mg atoms


**Q.2 (C)** The term **molecular formula cannot** be used for ionic compounds because molecular formula term is used for covalent compounds. In fact, molecule is an aggregation of atoms whereas **ionic compounds involve ions not atoms**.

**Q.3 (D)** Given amount of  $H_2SO_4 = 98g$

Number of moles of  $H_2SO_4 = \frac{98}{98}$

= 1mole

$H_2SO_4$  on dissociation splits up into ions such as



= 2 + 1 = 3moles of ions

**Conclusion:** From the equation it is clear that 1mole of  $H_2SO_4$  produces 3 moles of ions.

**Q.4 (D)** Cationic molecular ions can be generated by passing high energy beam of electrons,  $\alpha$ -particles or X-rays through a gas.

**Q.5 (A)** All the isotopes of an element have same number of protons and electrons but they have different mass number. e.g. Cl element has two isotopes  $^{35}_{17}Cl, ^{37}_{17}Cl$

**Q.6 (D)** A limiting reactant is the one which gives the minimum amount of the product under consideration. It is consumed earlier in the reaction.

**Q.7 (D)** Symbol does not represent amu.

$$1 \text{ amu} = \frac{1}{6.026 \times 10^{23}} \text{ g} = 1.661 \times 10^{-24} \text{ g}$$

$$\therefore 1 \text{ amu} = 1.661 \times 10^{-24} \text{ g}$$

$$= 1.661 \times 10^{-27} \text{ kg}$$

$$= 1.661 \times 10^{-21} \text{ mg}$$

**Q.8 (B)**

Opt.	Elements	No. of isotopes
A)	F	Mono-isotopic
B)	Cl	Di-isotopic
C)	Au	Mono-isotopic
D)	As	Mono-isotopic

**Q.9 (D)** Because cationic molecular ions are comparatively more stable than anionic molecular ions.

**Q.10 (A)**  $2C_2H_2 + 5O_2 \longrightarrow 4CO_2 + 2H_2O \dots$

Volume ratio b/w  $C_2H_2$  and  $O_2 = 2:5$

$2\text{cm}^3$  of  $C_2H_2$  requires  $O_2 = 5\text{cm}^3$

$1\text{cm}^3 \dots \dots \dots = \frac{5}{2}$

$5\text{cm}^3 \dots \dots \dots = \frac{5}{2} \times 5 = 12.5 \text{ cm}^3$

Total volume of oxygen gas required for complete combustion of ethyne =  $12.5\text{cm}^3$

**Q.11 (B)**  $B = 10.8$  amu (relative atomic mass of boron)

$$\begin{array}{l} \begin{array}{cc} & \swarrow \quad \searrow \\ {}^{11}\text{B} & & {}^{10}\text{B} \\ x & & 100-x \end{array} \\ \frac{11(x) + 10(100-x)}{100} = 10.8 \\ 11x + 1000 - 10x = 10.8 \times 100 \\ x + 1000 = 1080 \\ x = 1080 - 1000 = x = 80\% \end{array}$$

$\therefore$  %age of  ${}^{10}\text{B} = 100 - 80 = 20\%$

**Q.12 (C)** Number of C-atoms in sucrose

$$\begin{aligned} &= \frac{34.2}{342} \times 6 \times 10^{23} \times 12 \\ &= 7.2 \times 10^{23} \end{aligned}$$

**Q.13 (A)** Given data

Volume of nitrogen gas at

$$\text{RTP} = 1000\text{cm}^3$$

Number of nitrogen molecules ( $\text{N}_2$ )

$$\begin{aligned} &= \frac{1000}{24000} \times 6 \times 10^{23} \\ &= 2.5 \times 10^{22} \end{aligned}$$

**Q.14 (A)** Efficiency of a chemical reaction depends upon greater amount of product.

**Q.15 (A)** Total numbers of atoms in  $\text{H}_2\text{SO}_4$

$$\begin{aligned} &= \frac{49}{98} \times 6 \times 10^{23} \times 7 \\ &= 7 \times 3 \times 10^{23} \end{aligned}$$

**Q.16 (D)** Given data:

Empirical formula mass of organic compound ( $\text{CH}_2\text{O} = 30\text{g}$ )

Molecular mass of organic compound =  $90\text{g}$

Molecular formula of organic compound =  $n$  (Empirical formula)


$$n = \frac{\text{molecular mass}}{\text{empirical formula mass}} = \frac{90}{30} = 3$$

**Molecular formula** =  $3(\text{CH}_2\text{O}) = \text{C}_3\text{H}_6\text{O}_3$

**Q.17 (D)**

$$\text{Number of Br-atoms} = 3 \times 6.022 \times 10^{23}$$

**Q.18 (A)** When  $\text{CO}_2$  is absorbed in pre-weighed 50% KOH solution, reaction, take place as shown below:



From this reaction, it is clear that the absorption of  $\text{CO}_2$  in KOH solution is a **chemical change**.

**Q.19 (D)**  $\text{Mg}(\text{ClO}_4)_2$  acts as drying agent and absorbs water. Conc.  $\text{H}_2\text{SO}_4$  and  $\text{CaO}$  also act as drying agent.

**Q.20 (D)**  $\text{Mg} + \text{H}_2\text{SO}_4 \longrightarrow \text{MgSO}_4 + \text{H}_2$


$$\begin{array}{l} \text{Mg} : \text{H}_2 \\ 1 : 1 \\ 0.5 : 0.5 \text{ mole} \end{array}$$

- Number of moles of  $\text{Mg} = \frac{12}{24} = 0.5$
- Amount of  $\text{H}_2$  gas = Number of moles of  $\text{H}_2 \times$  molar mass of hydrogen gas  
 $0.5 \times 2 = 1.0\text{g}$
- Amount of  $\text{H}_2 = 1.0\text{g}$

**Q.21 (C)** The percentage oxygen is obtained by the method of difference

$$\% \text{ of O} = 100 - (\% \text{ of C} + \% \text{ of H})$$

**Q.22 (B)**



$$0.36 \quad 0.36 \quad ?$$

(a) (b)

(a)  $\text{Al} : \text{Al}_2\text{O}_3$

$$4 : 2$$

$$0.36 : x$$

$$x = \frac{0.36 \times 2}{4} = 0.18$$

(b)  $\text{O}_2 : \text{Al}_2\text{O}_3$


$$3 : 2$$

$$0.36 : x$$

$$x = \frac{0.36 \times 2}{3} = 0.24$$

Conclusion by comparison of (a) and (b) it is clear that aluminum produces least amount of

product. So, aluminum acts as limiting reactant while oxygen acts as non-limiting reactant.



Mass of sand ( $\text{SiO}_2$ ) is treated with

$$\text{C} = 0.3\text{kg} = 300\text{g}$$

Mass of Silicon carbide produced (actual yield)

$$= 0.1\text{kg} = 100\text{g} \dots \text{i}$$

Molar mass of sand ( $\text{SiO}_2$ ) = 28 + 32

$$= 60.0\text{gmol}^{-1}$$

Molar mass of silicon carbide = 28 + 12

$$= 40\text{gmol}^{-1}$$

$$\text{Theoretical Yield} = \frac{40}{60} \times 300 = 200\text{g} \dots$$

ii

$$\text{Percentage Yield} = \frac{\text{Actual Yield}}{\text{Theoretical Yield}} \times 100$$

$$= \frac{100}{200} \times 100 = 50\% \dots \text{iii}$$

$\therefore$  Percentage Yield of silicon carbide ( $\text{SiC}$ ) = 50%

**Q.24 (B)**

- Relative isotopic mass term is used only for isotopes
- Moreover, carbon element has three isotopes  ${}^{12}_6\text{C}$ ,  ${}^{13}_6\text{C}$ ,  ${}^{14}_6\text{C}$  but not  ${}^{15}_{16}\text{C}$

**Q.25 (D)** In fact, the term mass number is used for isotopes of an element. Mass number is sum of protons and neutrons but it is not sum of protons and electrons.

**Q.26 (C)** In fact, both experimental error and human error affect actual yield.

**Q.27 (A)** Since all the isotopes of an element have same proton number, therefore, they have same electronic configuration. So isotopes of an element have same chemical properties but have different physical properties because they have different mass numbers.

**Q.28 (A) Given data**

C %	:	H %	:	O %
60	:	8.0	:	32
$\frac{60}{12}$	:	$\frac{8.0}{1}$	:	$\frac{32}{16}$
5	:	8	:	2
2.5	:	4	:	1
2(2.5	:	4	:	1)
Empirical formula				= $\text{C}_5\text{H}_8\text{O}_2$
Empirical formula mass				= 100
Molecular formula				= n(empirical formula)
				$n = \frac{200}{100} = 2$

$\therefore$  Molecular formula =  $\text{C}_{10}\text{H}_{16}\text{O}_4$

**Q.29 (D)**

C %	:	H %	:	O %
48	:	4	:	48
$\frac{48}{12}$	:	$\frac{4}{1}$	:	$\frac{48}{16}$
4	:	4	:	3
3	:	3	:	3
3(1.33	:	1.33	:	1)
4	:	4	:	3

Empirical Formula of ascorbic acid =  $\text{C}_4\text{H}_4\text{O}_3$

**Q.30 (B) From the balanced equation**



$$\text{Molar ratio } 2 : 2$$

$$0.1 : 0.1$$

$$\text{Molar mass of CaO} = 56\text{amu}$$

$$\text{Mass of CaO formed} = 0.1 \times 56 = 5.6\text{g}$$

**Q.31 (B)** If we know the mass of one substance, we can calculate the volume of other substance with the help of balanced chemical equation and this relationship is called mass – volume relationship.

**Q.32 (A)** Number of chlorine atoms

$$= \frac{7.1}{35.5} \times 6 \times 10^{23}$$

$$= 1.2 \times 10^{23}$$

## Q.33 (B)

	CO <sub>2</sub>	:	H <sub>2</sub> O
Molar mass	44g	:	18g
According to Condition	11g	:	?
Amount of water			= 4.5g

∴ 4.5g of water has same number of water molecule as present in 11g of CO<sub>2</sub>

Q.34 (A) 1mole of N<sub>2</sub> gas = 28g  
= 22.41dm<sup>3</sup> at STP

Q.35 (D) Both A and B

- (A) 12.0 g of <sup>12</sup>C contains 1 mole or  $6.02 \times 10^{23}$  carbon atoms.  
 (B) 4.0 g of <sup>4</sup>He also contains 1 mole of He atoms.  
 (C) 2.0 g of <sup>1</sup>H<sub>2</sub> contains 1 mole of H<sub>2</sub> molecules but 2 moles of H atoms.




Q.36 (D) Let X be the number of C atoms per molecule of X. therefore, 1 mole of X will produce X moles of CO<sub>2</sub> when completely burnt in O<sub>2</sub>.

$$\begin{aligned} \text{Amount of x used} &= \frac{0.112}{22.4} \\ &= 5.00 \times 10^{-3} \text{ mol} \end{aligned}$$

$$\begin{aligned} \text{Amount of CO}_2 \text{ produced} &= \frac{0.88}{12 + 2(16)} = \frac{0.88}{44} \\ &= 2.0 \times 10^{-2} \text{ mol} \end{aligned}$$

$$\begin{aligned} \therefore (5.00 \times 10^{-3})x &= 2.0 \times 10^{-2} \quad x = \frac{2.0 \times 10^{-2}}{5.00 \times 10^{-3}} \\ x &= 4 \text{ Ans.} \end{aligned}$$

Q.37




## Q.38 (C)

	C	H	O
% mass	72	12	16
Relative atomic mass	12	1	16
Molar ratio	$\frac{72}{12}$	$\frac{12}{1}$	$\frac{16}{16}$
	6	12	1
Empirical formula	C <sub>6</sub> H <sub>12</sub> O		

Q.39 (D) The isotopes of Cl are <sup>35</sup>Cl and <sup>37</sup>Cl and they exist in the ratio of 3 : 1.  
Hence

$$\begin{aligned} \text{A. of Cl} &= \left( \frac{3}{1+3} \times 35 \right) + \left( \frac{1}{1+3} \times 37 \right) \\ &= 35.5 \text{ amu} \end{aligned}$$

The mass spectrograph of chlorine element is:



Q.40 (B)

$$\begin{aligned} \text{Molar mass of MgSO}_4 &= 1 \times 24 + 1 \times 32 + 4 \times 16 \\ &= 24 + 32 + 64 = 120\text{g} \end{aligned}$$

$$\begin{aligned} \text{Mass of MgSO}_4 &= \text{mole} \times \text{molar mass of MgSO}_4 \\ &= 10^{-3} \times 120 \\ &= 0.12 \text{ g} \end{aligned}$$

# STOP

A PROGRAM BY PUNJAB GROUP

